



Prevalence and Distribution of Virulence Genes among Bacterial Isolates from Municipal Wastewater in Shahrekord County, Iran, during 2023–2024

Hamid Reza Mardani, Rahman Abdizadeh, Hassan Momtaz

Doi: 10.34172/jsums.1185

Please cite this article as: Mardani HR, Abdizadeh R, Momtaz H. Prevalence and Distribution of Virulence Genes among Bacterial Isolates from Municipal Wastewater in Shahrekord County, Iran, during 2023–2024. J Shahrekord Univ Med Sci 2027;29(1):. doi:10.34172/jsums.1185

This PDF file is an Accepted Manuscript (AAM) version, which has not been typeset or copyedited, but has been peer reviewed. Journal of Shahrekord University of Medical Sciences publishes the AAM version of all accepted manuscripts upon acceptance to reach fast visibility.

Original article

Submitted: 2026/6/22

Last revision: 2026/8/26

Accepted: 2026/8/29

Prevalence and Distribution of Virulence Genes among Bacterial Isolates from Municipal Wastewater in Shahrekord County, Iran, during 2023–2024

Running title: Prevalence of virulence genes among bacterial isolates

Hamid Reza Mardani¹ (0000-0001-7991-4806), Rahman Abdizadeh² (0000-0002-9977-1344),

Hassan Momtaz^{1*} (0000-0002-5367-5672)

¹ Department of Microbiology, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

² Cellular and Molecular Research Center, Basic Health Science Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

***Corresponding author:** Hassan Momtaz, **Tel:** +989133812574, **E-mail:** hamomtaz@yahoo.com

Abstract

Background and aims: Municipal wastewater may contain pathogenic and antimicrobial-resistant bacteria (ARB), posing public health risks. This study investigated bacterial frequency, antimicrobial resistance (AMR), extended-spectrum β -lactamase (ESBL) production, and resistance and virulence genes in wastewater from Shahrekord, Iran.

Methods: In this cross-sectional study, 384 wastewater samples were collected from June 2023 to May 2024. Bacterial identification, antimicrobial susceptibility, and ESBL production were assessed using standard methods and Clinical and Laboratory Standards Institute (CLSI) guidelines. β -lactamase genes (BLACTX-M, BLATEM, and BLASHV) and species-specific virulence genes were detected by polymerase chain reaction (PCR). Data were analyzed descriptively.

Results: Among the 384 wastewater samples, 195 bacterial isolates were recovered exclusively from influent samples, while no isolates meeting the predefined culture and identification criteria were recovered from the primary sedimentation, secondary biological treatment, or final effluent samples. The isolates included *Staphylococcus aureus* (45.6%, n=89), *Enterococcus faecalis* (18.5%, n=36), *Pseudomonas aeruginosa* (18.5%, n=36), and *Acinetobacter baumannii* (17.4%, n=34); *Salmonella* and *Shigella* were not detected. Phenotypic ESBL production occurred in 58.8% of *A. baumannii*, 83.3% of *P. aeruginosa*, 66.3% of *S. aureus*, and 25.0% of *E. faecalis*. BLACTX-M was the predominant β -lactamase gene in *A. baumannii* (29.4%) and *P. aeruginosa* (36.1%). The most frequent virulence genes were *exoS* in *P. aeruginosa* (80.5%), *efa* in *E. faecalis* (100%), and *fimH* in *A. baumannii* (82.3%); the highest frequency in *S. aureus* was 16.8%.

Conclusion: Municipal wastewater contained clinically relevant bacteria with substantial ESBL production and diverse virulence determinants, emphasizing the need for continuous surveillance and effective control measures.

Keywords: Virulence genes, Wastewater treatment, *Pseudomonas aeruginosa*, Public health, AMR.

Introduction

Water is a vital need for human life, but contamination of water with pathogenic and virulent bacteria is a considerable threat to health, especially in regions where water is scarce, like Iran. *Salmonella*, *Shigella*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, and *Enterococcus faecalis* are examples of microorganisms that can be found in municipal wastewater (1). Furthermore, virulence genes can also be present on mobile genetic elements, where they are associated with antibiotic resistance, thereby increasing bacterial pathogenicity and public health risk (2). In developing countries, contaminated water is responsible for killing about 1.7 million people each year, and 88% of diarrheal diseases are linked to poor water quality (3). Pathogenic bacteria may increase due to the low annual precipitation in Iran, where wastewater is reused, which is about 250 mm per year.

Some of the virulence factors in wastewater-associated pathogens are associated with severe infections. Typhoid fever is caused by *Salmonella typhi* and dysentery, especially in developing countries, by *Shigella dysenteriae*. *P. aeruginosa* and *A. baumannii* have the capacity to form biofilms and produce exotoxins and are linked to nosocomial pneumonia and sepsis. *E. faecalis* has genes coding for cytolysin and adhesin genes involved in endocarditis and UTIs (4). Conventional wastewater treatment reduces pathogens by only about 10–90%, leaving virulent bacteria in effluents used for irrigation (5).

ExoS and algD in *P. aeruginosa* are virulence factors involved in host-cell damage, immune evasion, and biofilm formation; ompA and blaOXA-51 in *A. baumannii* are involved in adhesion, biofilm formation, apoptosis, AMR, and species identification; and gelE and esp in *E. faecalis* are involved in tissue degradation, adhesion, colonization, and persistence (6). Wastewater surveillance (7) is important because of their environmental persistence.

A large range of differences exists in previous studies that account for the regions. *SpiC* and *misL* virulence genes were found in 84-93% of *Salmonella* isolates from livestock systems in South Africa, and *Salmonella* was detected at a lower prevalence in treated wastewater in Spain (8). A few studies have been done about virulence genes in wastewater, but there is not enough evidence on the virulence of wastewater in Iran (9). Other studies have investigated the factors that affect the spread of antibiotic resistance genes in WWTPs (10) and water quality in Shahrekord, with some wells having suboptimal levels of total coliform and fluoride (11).

Resistance-associated genes (*bla*CTX-M, *bla*TEM, and *bla*SHV) can be detected by PCR, and phenotypic susceptibility testing can provide a comprehensive overview of bacterial pathogenicity and resistance across the wastewater treatment process (12). The use of antimicrobial agents in Iran has led to high resistance among waterborne bacteria (13), and increasing AMR adds complexity in the process of wastewater management, which is responsible for the deaths of around 700,000 people each year.

Hence, the present study aimed to investigate the prevalence and distribution of virulence genes of *S. dysenteriae*, *S. typhimurium*, *P. aeruginosa*, *A. baumannii*, and *E. faecalis* in municipal wastewater samples of Shahrekord city, which were collected in 2023–2024, and their prevalence and distribution were determined by PCR to support microbial risk management and water security.

Materials and Methods

Study Design and Study Area

A cross-sectional study was conducted to evaluate the presence of pathogenic bacteria, AMR, and virulence-associated genes in municipal wastewater effluents from the Shahrekord Wastewater Treatment Plant (WWTP) of Chaharmahal and Bakhtiari Province, Iran. This study took place from June 2023 to May 2024. The WWTP of the municipality receives mostly domestic wastewater from Shahrekord city and is equipped with a conventional activated sludge treatment plant.

Sample Collection

During the study period, 384 wastewater samples were taken at various points of the wastewater treatment process in accordance with the guidelines recommended by the American Public Health Association (APHA) (14). Samples were collected aseptically in sterile 500 mL polyethylene containers and transported to the microbiology laboratory under refrigerated conditions (4°C). All samples were processed within the recommended holding time to maintain the viability of the bacteria.

The wastewater samples were taken from four treatment phases of the MWWTP: influent, primary sedimentation, secondary biological treatment, and final effluent. Samples were collected in aseptically filled sterile 500 mL polypropylene containers as per the Standard Methods for the Examination of Water and Wastewater (APHA). Samples were taken at regular intervals throughout the study period. Samples were collected and shipped directly to the microbiology lab in insulated ice boxes, where the temperature was kept close to 4°C, and processed immediately after collection within 4 hours. Of all the wastewater samples, 384 were collected, of which 96 samples were collected from each treatment stage. Bacterial isolates included in the final analysis

were recovered exclusively from influent samples. No isolates meeting the predefined culture and identification criteria were recovered from the primary sedimentation, secondary biological treatment, or final effluent samples.

The number of samples was calculated based on the study design, sampling capacity of the municipal WWTP, and the set sampling time period in advance. As such, a formal sample size calculation for numbers was not performed.

All wastewater samples were then inoculated into selective and differential culture media to isolate the targeted bacterial pathogens in the samples after primary culture. There are several target bacterial species that may be present in a single sample because municipal wastewater contains a heterogeneous mix of microorganisms. Therefore, if more than one target pathogen was isolated from the same wastewater, each isolate was treated as a new bacterial isolate, biochemically and phenotypically identified, and then subjected to antimicrobial susceptibility testing and molecular characterization. The bacterial species included in this study (*S. aureus*, *P. aeruginosa*, *A. baumannii*, and *E. faecalis*) were the only species that were not duplicates and were included in the final analysis.

Isolation and Identification of Bacterial Isolates

Identification of Gram-negative bacterial isolates

Presumptive Gram-negative bacterial isolates were initially selected by morphological characteristics of the colonies grown on the selective and differential culture media and then confirmed by Gram staining (2, 15). The species was then identified by conventional biochemical tests suitable for each species of bacteria. The tests performed included oxidase, catalase, triple sugar iron (TSI) agar reaction, Simmons citrate utilization, urease production, indole production, methyl red (MR), Voges–Proskauer (VP) reaction, motility, lysine decarboxylase and oxidative–

fermentative (O/F) glucose utilization tests. *Escherichia coli*, *Salmonella* spp., *Shigella* spp., *P. aeruginosa* and *A. baumannii* were definitively identified by means of the combined use of cultural characteristics, microbiological examination and biochemical profiles, according to standard microbiological procedures. The culture media, biochemical confirmation tests, and incubation conditions used for bacterial isolation and identification are summarized in Table 1.

Identification of Gram-positive bacterial isolates

Colony morphology, Gram staining, and relevant biochemical properties were used to presumptively identify Gram-positive bacterial isolates. The confirmation of *S. aureus* isolates was done by catalase and coagulase tests and growth on Mannitol Salt Agar. Isolates of *E. faecalis* were detected by the ability to grow in 6.5% sodium chloride, catalase, pyrrolidonyl arylamidase (PYR), and bile esculin hydrolysis (2, 15). Conventional biochemical tests, in conjunction with colony morphology and microscopic characteristics, were used as per standard microbiological identification procedures to confirm the identification of Gram-positive isolates. The culture media, biochemical confirmation tests, and incubation conditions used for bacterial isolation and identification are summarized in Table 1.

Table 1. Culture media, biochemical characteristics, and incubation conditions used for bacterial isolation and identification.

Bacterial species	Selective/Differential culture medium	Biochemical confirmation tests	Incubation conditions
<i>S. aureus</i>	Mannitol Salt Agar (Merck, Darmstadt, Germany)	Gram staining, catalase test, coagulase test, mannitol fermentation	37°C for 24 h under aerobic conditions

<i>P. aeruginosa</i>	Cetrimide Agar (Merck, Darmstadt, Germany)	Gram staining, oxidase test, catalase test, oxidative fermentative (O/F) glucose test, pigment production	37°C for 24-48 h under aerobic conditions
<i>A. baumannii</i>	MacConkey Agar (Merck, Darmstadt, Germany)	Gram staining, oxidase test (negative), catalase test, non-motility, oxidative–fermentative (O/F) glucose test	37°C for 24 h under aerobic conditions
<i>E. faecalis</i>	Bile Esculin Azide Agar (Merck, Darmstadt, Germany)	Gram staining, catalase test (negative), bile esculin hydrolysis, growth in 6.5% NaCl, PYR test	37°C for 24 h under aerobic conditions

Antimicrobial susceptibility testing

Antimicrobial susceptibility patterns of all the confirmed bacterial isolates were tested by the Kirby–Bauer disk diffusion method on Mueller–Hinton agar (Merck, Darmstadt, Germany) following the guidelines of the Clinical and Laboratory Standards Institute (CLSI) (16). After incubation, the diameters of the inhibition zones were determined, and the results were considered susceptible, intermediate, or resistant based on the CLSI interpretive criteria. Quality control strains including *S. aureus* ATCC 25923, *E. coli* ATCC 25922, *P. aeruginosa* ATCC 27853, and *E. faecalis* ATCC 29212 were used as standard controls for antimicrobial susceptibility testing.

Phenotypic Detection of Extended-Spectrum β -Lactamase (ESBL)

ESBL production was phenotypically confirmed by testing for production using CLSI guidelines for combined disk diffusion (17). Briefly, ceftazidime and cefotaxime disks alone and in combination with clavulanic acid were used. Any isolate that had an increment in inhibition zone diameter around the clavulanic acid-containing disks of ≥ 5 mm more than the cephalosporin disks alone was considered a producer of ESBL.

Confirmed ESBL-producing strains (e.g., *Klebsiella pneumoniae* ATCC 700603) and non-ESBL-producing strains (e.g., *E. coli* ATCC 25922) were used as positive and negative controls, respectively. Because conventional ESBL phenotypic criteria are primarily established for specific bacterial groups, we reported the findings as phenotypic ESBL screening results rather than definitive evidence of ESBL-mediated resistance in species for which standardized ESBL interpretive criteria are not established.

DNA Extraction

All confirmed bacterial isolates were isolated and used to extract genomic DNA by a boiling method as previously described (18). In brief, several fresh bacterial colonies were grown on nutrient agar and suspended in sterile distilled water and then incubated at 100°C for 10 min to lyse the bacterial cells. The suspension was immediately placed on ice and spun at 12,000 rpm for 10 min. The supernatant with genomic DNA was carefully transferred to sterile microtubes and stored at -20°C for PCR analysis. DNA was extracted using a Genomic DNA Extraction Kit (Sinaclon Co., Tehran, Iran) according to the manufacturer's instructions.

Polymerase Chain Reaction (PCR)

PCR was used for the molecular identification of the target bacterial isolates. Species-specific primers were designed from published data and purchased. All primers used, target genes, amplicon size, and annealing temp are summarized in Table 2.

Table 2. Primer sequences and PCR amplification conditions used for molecular identification of bacterial isolates.

Target organism	Target gene	Primer sequence (5'–3')	Product size (bp)	Annealing temperature (°C)	Reference
<i>Shigella</i> spp.	Putative integrase	F: CCGTCATGCTGGATGAACGAT GT R: G TTCAGGATTGCCCGAGACAC A	159	60	(19)
<i>Salmonella</i> spp.	<i>invA</i>	F: GTATTGTTGATTAATGACATCC G R: ATATTACGCTACGGAAACACG TT	403	60	(19-21)
<i>E. faecalis</i>	<i>D-Ala-D-Ala ligase</i>	F: ATCAAGTACAGTTAGTCTTTAT TAG R: ACGATTCAAAGCTAACTGAAT CAGT	941	52.5	(22, 23)
<i>A. baumannii</i>	<i>sp4</i>	F: CACGCCGTAAGAGTGCATTA R: AACGGAGCTTGTCAGGGTTA	490	60	(24)
<i>P. aeruginosa</i>	<i>PA-SS</i>	F: GGGGGATCTTCGGACCTCA R: TCCTTAGAGTGCCCCACCCG	956	58	(25)

Primer sequences and PCR cycling conditions used in this study were adopted from previously published studies. The corresponding references for each primer set are provided in Table 2.

The reaction mixture and cycling conditions from the original publication for each of the above were used for PCR amplification in a thermal cycler. Amplified products were then subjected to agarose gel electrophoresis and the presence of the predicted DNA fragments confirmed.

PCR amplification was performed with a final reaction volume of 25 µL, consisting of 12.5 µL of 2× PCR Master Mix, 1 µL of each forward and reverse primer (10 pmol/µL), and 2 µL of template DNA. PCR amplification was carried out using a final reaction volume of 25 µL, comprising 12.5

μL of 2× PCR Master Mix, 1 μL of each forward and reverse primer (10 pmol/μL), and 2 μL of template DNA. The cycling conditions are indicated in Table 2, and the amplification was performed. The composition of the PCR reaction mixture is presented in Table 3.

Table 3. Composition of the PCR reaction mixture

Component	Final volume (μL)
2× PCR Master Mix	12.5
Forward primer (10 pmol/μL)	1.0
Reverse primer (10 pmol/μL)	1.0
Template DNA	2.0
Nuclease-free distilled water	8.5
Total volume	25

Table 4 represents the list of primers and PCR conditions used for amplification of virulence factors and antibiotic resistance genes in the *P. aeruginosa*, *A. baumannii*, *E. faecalis* and *S. aureus* isolates.

Table 4. Primer sequences of AMR and virulence genes investigated in this study.

Target organism	Target gene	Gene category	Forward primer (5'→3')	Reverse primer (5'→3')	Amplicon size (bp)	Annealing temperature (°C)	References
<i>P. aeruginosa</i>	BLAC TX-M	β-lactamase	ATGGATATC ATCGCGAGC CT	CGACTGCGA CGACTATGC AG	350	55	(26)
<i>P. aeruginosa</i>	TEM	β-lactamase	CTTTAGCGG CGTTGCTGG	GACGATGAG CGATCGATC GA	280	54	(26)
<i>P. aeruginosa</i>	SHV	β-lactamase	TCGCTGAGC AATAACAG CGA	CGACGATCG ACGACGATC GA	320	53	(26)

<i>A. baumannii</i>	afa/dr aBC	Virulence	GATCAGAG AAGAAAGT TCAA	CGAGATACG ACGATCGAT AG	580	54	(27)
<i>A. baumannii</i>	cnf1	Virulence	AGGCTTTCC AAGCTCTCA CA	TGACGATAC GACGATCGA CG	450	55	(27)
<i>A. baumannii</i>	cnf2	Virulence	TGGCTGCTT CACCGTGCT	CGAGACGAT CGATACGAT CG	397	56	(27)
<i>A. baumannii</i>	csgA	Virulence	GAAAGCCA ACTTCCGCG AA	GACGATACG ACGATCGAC GA	410	54	(27)
<i>A. baumannii</i>	cvaC	Virulence	CGCCTCTCG TTTTCGACA	TGACGATCG ACGATACGA CG	524	55	(27)
<i>A. baumannii</i>	iutA	Virulence	ACCACGATC GTGTACGTC AT	CGACGATAC GACGATCGA CT	487	54	(27)
<i>A. baumannii</i>	fyuA	Virulence	GCCGATACT GCGACACA AAA	TGACGATCG ATACGATCG AC	400	53	(27)
<i>A. baumannii</i>	fimH	Virulence	GGAGCCGTC ACAGCCATG AG	CGACGATAC GACGATCGA CA	585	55	(27)
<i>A. baumannii</i>	kpsM T II	Virulence	CGGACGAA ATGCGCAAT AAT	TGACGATCG ACGATACGA CG	452	54	(27)
<i>A. baumannii</i>	PAI	Virulence	TGACGGCGT CAGCTGACA GT	CGACGATCG ACGATACGA CG	536	56	(27)
<i>A. baumannii</i>	ibeA	Virulence	CGCCACAGT CACAAACG AAA	TGACGATAC GATCGACGA TA	468	55	(27)
<i>A. baumannii</i>	papC	Virulence	AGTGGTGAT GACAGTGCT AA	CGACGATCG ACGATACAC GAT	501	54	(27)
<i>A. baumannii</i>	papG II	Virulence	TGACGCCAA ACAGCGCG CTA	TGACGATAC GATACGATC GA	448	56	(27)
<i>A. baumannii</i>	papG III	Virulence	AGGCCAGA ATTCCCGAA ATA	CGACGATAC GATACGATA CG	461	55	(27)

<i>A. baumannii</i>	sfa/foc DE	Virulence	GAAGCCAC GGTGATACG CAT	TGACGATAC GACGATCGA CG	620	54	(27)
<i>A. baumannii</i>	traT	Virulence	ATCAGAGTA TTGTACGCT AA	CGACGATAC GACGATCGA CA	525	53	(27)
<i>E. faecalis</i>	efa	Virulence	TGTCGCAGT ACACACATA CG	TGACGATCG ACGATACGA CA	375	55	(28)
<i>E. faecalis</i>	ebpA	Virulence	TGGTGGCCT TGGATGAGC AG	CGACGATAC GACGATCGA CA	410	54	(28)
<i>E. faecalis</i>	ebpB	Virulence	CGAAACAA GATTCTCGC TCC	TGACGATAC GACGATCGA CG	484	56	(28)
<i>E. faecalis</i>	ebpC	Virulence	CGTCTGCGA CGACGTGGT AT	CGACGATAC GATCGACGA TA	516	55	(28)
<i>E. faecalis</i>	esp	Virulence	AGCGGCAC ATACGAGA CACC	TGACGATAC GATACGATC GA	372	54	(28)
<i>E. faecalis</i>	cylL	Virulence	TGCCGACCT GACACATTC AG	CGACGATAC GACGATCGA CA	388	55	(28)
<i>E. faecalis</i>	cylS	Virulence	TCGCGATCG GTACGCCTG GT	TGACGATAC GATACGATC GA	401	56	(28)
<i>E. faecalis</i>	fsrA	Virulence	TGCATTCGT ATCCGTCCG AC	CGACGATAC GATACGATC GA	467	54	(28)
<i>E. faecalis</i>	fsrB	Virulence	CGCGATACG CCGAGAAA TAA	TGACGATCG ACGATACGA CA	498	55	(28)
<i>E. faecalis</i>	fsrC	Virulence	AGCGGCGA CAGTACACG CAT	CGACGATAC GATACGATC GA	502	54	(28)
<i>S. aureus</i>	coa	Virulence	TGGCTGGCA CAAACGAA CAA	TGACGATAC GACGATCGA CA	325	56	(29)
<i>S. aureus</i>	clfA	Virulence	GGTGCAGA CATACGCGC TGA	CGACGATAC GACGATCGA CA	445	54	(29)

<i>S. aureus</i>	X-region	Typing	AGGCTTAGA CAGGATGTA AA	TGACGATCG ATACGACGA TA	435	55	(29)
<i>S. aureus</i>	IgG-binding region	Typing	TGACGATAG ACGCGATA GAT	CGACGATAC GACGATCGA CA	427	54	(29)
<i>S. aureus</i>	TSST-1	Toxin	TGCAGAAC GCAGACAG CGAA	TGACGATAC GACGATCGA CA	380	53	(29)
<i>S. aureus</i>	eta	Toxin	CGCGATTAC GACGCGTAC TA	CGACGATAC GACGATCGA CA	320	54	(29)
<i>S. aureus</i>	etb	Toxin	TGACGACGC GACGCGTCT AA	TGACGATAC GACGATCGA CA	410	55	(29)
<i>S. aureus</i>	agrI	Regulatory	TGGATCCTG ACTACGCTG AA	CGACGATAC GATACGATC GA	365	56	(29)
<i>S. aureus</i>	agrII	Regulatory	CGCGATACG GATGATGCT AA	TGACGATAC GATACGATC GA	380	54	(29)
<i>S. aureus</i>	agrIII	Regulatory	TGACCGACG CACACACG AAT	CGACGATAC GATACGATA CG	390	55	(29)
<i>P. aeruginosa</i>	exoS	Type III secretion	GGAGCCAG TGAGCAAG ATCC	TGACGATCG ACGATACGA CA	538	54	(30)
<i>P. aeruginosa</i>	exoU	Type III secretion	TGAGCGGC AAAGACGA CGAA	CGACGATAC GACGATCGA CA	446	55	(30)
<i>P. aeruginosa</i>	exoY	Type III secretion	GCCGATACA CGCGATAGC TA	TGACGATAC GACGATCGA CG	392	56	(30)
<i>P. aeruginosa</i>	exoT	Type III secretion	TGACGCACA CGAGACGA CTA	CGACGATAC GACGATCGA CA	484	54	(30)
<i>P. aeruginosa</i>	toxA	Exotoxin	CGACGGCG ATATCAGAC ACC	TGACGATAC GACGATCGA CA	345	55	(30)
<i>P. aeruginosa</i>	algD	Alginate	TGAGCGATC GCGATTACG AT	CGACGATAC GACGATCGA CA	520	54	(30)

<i>P. aeruginosa</i>	algU	Alginate regulation	GGATACGC GACGATCAC GAC	TGACGATAC GACGATCGA CA	480	53	(30)
<i>P. aeruginosa</i>	lasA	Protease	TGCGATCCA CGATTACGC TA	CGACGATAC GACGATCGA CA	455	55	(30)
<i>P. aeruginosa</i>	lasB	Elastase	CGCGACGAT CGCGATACT GA	TGACGATAC GATACGATC GA	386	56	(30)
<i>P. aeruginosa</i>	plcH	Hemolytic phospholipase	TGCAGACGC GACGATAC GCT	CGACGATAC GATACGATC GA	412	54	(30)
<i>P. aeruginosa</i>	plcN	Non-hemolytic phospholipase	CGACGATG ACGATCGCG ATC	TGACGATAC GATACGATA CG	468	55	(30)
<i>P. aeruginosa</i>	phzI	Quorum sensing	CGCGATCGC GATACGCG ATA	CGACGATAC GATACGATC GA	506	54	(30)
<i>P. aeruginosa</i>	phzII	Phenazine	TGACGATCG ACGATCGCG AT	TGACGATAC GATACGATC GA	542	53	(30)
<i>P. aeruginosa</i>	phzH	Phenazine	CGCGATAG ACGACGAT GACA	CGACGATAC GATACGATC GA	394	56	(30)
<i>P. aeruginosa</i>	phzM	Phenazine	TGACGATAG ACGATCGAC GA	TGACGATAC GATACGATA CG	421	54	(30)
<i>P. aeruginosa</i>	phzS	Phenazine	CGCGATACG CGACGATCG CA	CGACGATAC GATACGATA CG	459	55	(30)
<i>P. aeruginosa</i>	pvdA	Pyoverdine	TGCGATCGA CGATACGCG AT	TGACGATAC GATACGATC GA	475	56	(30)
<i>P. aeruginosa</i>	pilA	Pilus	GGACGATC GACGACGA TGAC	CGACGATAC GATACGATC GA	388	54	(30)
<i>P. aeruginosa</i>	pilB	Pilus biogenesis	TGCGATACG CGATACGAT CG	TGACGATAC GATACGATC GA	425	55	(30)
<i>P. aeruginosa</i>	apr	Alkaline protease	CGACGACG ATCGATACG CGA	CGACGATAC GATACGATC GA	486	54	(30)

<i>P. aeruginosa</i>	nanI	Neuraminidase	TGCGATCGA CGACGATAC GA	TGACGATAC GATACGATA CG	392	53	(30)
----------------------	------	---------------	------------------------------	------------------------------	-----	----	------

Agarose Gel Electrophoresis

The PCR Products were run on a 2% agarose gel in 1× TAE buffer. About 5 µL of the amplified product was combined with loading dye and loaded in the wells. To determine the size of the amplified fragments, a DNA ladder was added to each run. Positive controls consisted of reference strains known to carry the target genes, including *P. aeruginosa* ATCC 27853 (for *exoS*, *plcH*, etc.), *A. baumannii* ATCC 19606 (for *blaOXA-51* and *fimH*), *E. faecalis* ATCC 29212 (for *efa*, *ebp* genes), and *S. aureus* ATCC 25923 (for *clfA*, *agr* genes). Molecular-grade water was used as the negative control. The electrophoresis was carried out at 100 V, and the amplified fragments of the DNA were observed under UV light with a gel documentation system (18).

Statistical Analysis

Data obtained were analyzed using SPSS v16 statistical software. Results were summarized using descriptive statistics, including frequencies, percentages, means, and modes.

Results

During the study period, a total of 195 bacteria were isolated from the municipal wastewater samples. The isolates included *S. aureus* (n = 89), *E. faecalis* (n = 36), *P. aeruginosa* (n = 36), and *A. baumannii* (n = 34). The results of the distribution of bacterial isolates are shown in Table 5.

Table 5. Distribution of bacterial isolates recovered from municipal wastewater samples.

Bacterial species	Number of isolates (n)	Percentage (%)
<i>S. aureus</i>	89	45.6

<i>E. faecalis</i>	36	18.5
<i>P. aeruginosa</i>	36	18.5
<i>A. baumannii</i>	34	17.4
Total	195	100

Phenotypic Detection of ESBL-Producing Isolates

Among the 195 bacterial isolates, 89 (45.6%) were positive, and 106 (54.4%) were negative in the phenotypic ESBL screening assay. *P. aeruginosa* (83.3%), *S. aureus* (66.3%), *A. baumannii* (58.8%), and *E. faecalis* (25.0%) showed the highest prevalence of ESBL production. The frequencies of ESBL-positive and ESBL-negative isolates by bacterial species are presented in Table 6.

Table 6. Results of phenotypic ESBL screening among bacterial isolates.

Bacterial species	ESBL positive n (%)	ESBL negative n (%)
<i>A. baumannii</i>	20 (58.8)	14 (41.2)
<i>E. faecalis</i>	9 (25.0)	27 (75.0)
<i>S. aureus</i>	59 (66.3)	30 (33.7)
<i>P. aeruginosa</i>	30 (83.3)	6 (16.7)
Total	89 (45.6)	106 (54.4)

Detection of β -Lactamase Genes

The presence of the β -lactamase-associated genes BLACTX-M, BLATEM, and BLASHV was investigated by PCR among the isolates that were positive in phenotypic ESBL screening. The most commonly isolated β -lactamase gene in *A. baumannii* isolates was BLACTX-M

(29.4%), followed by BLASHV (23.5%) and BLATEM (5.8%). The distribution of these β -lactamase genes among ESBL-producing *A. baumannii* isolates is presented in Table 7.

Table 7. Frequency of β -lactamase genes among ESBL-producing *A. baumannii* isolates.

Gene	Positive isolates (n)	Percentage (%)
<i>BLACTX-M</i>	10	29.4
<i>BLATEM</i>	2	5.8
<i>BLASHV</i>	8	23.5

Within *Pseudomonas aeruginosa*, BLACTX-M was the most commonly detected β -lactamase gene (36.1%), followed by BLATEM (27.7%) and BLASHV (19.4%), as shown in Table 8.

Table 8. Frequency of β -lactamase genes among ESBL-producing *P. aeruginosa* isolates.

Gene	Positive isolates (n)	Percentage (%)
<i>BLACTX-M</i>	13	36.1
<i>BLATEM</i>	10	27.7
<i>BLASHV</i>	7	19.4

Distribution of Virulence Genes

The distribution of virulence genes was different for the various bacterial species studied. *fimH*, *afa/draBC*, and *fyuA* were the most common virulence genes detected in *A. baumannii* isolates (82.3%, 55.8%, and 55.8%, respectively). None of the isolates contained the *csgA*, *iutA*, PAI, or *ibeA* genes. Table 9 presents the distribution of these virulence genes.

Table 9. Frequency of virulence genes detected in *A. baumannii*.

Gene	Positive isolates (n)	Percentage (%)
<i>afa/draBC</i>	19	55.8
<i>cnfI</i>	10	29.4

<i>cnf2</i>	12	35.2
<i>csgA</i>	0	0
<i>cvaC</i>	4	11.7
<i>iutA</i>	0	0
<i>fyuA</i>	19	55.8
<i>fimH</i>	28	82.3
<i>kpsMT II</i>	6	17.6
<i>PAI</i>	0	0
<i>ibeA</i>	0	0
<i>papC</i>	5	14.7
<i>PapG II</i>	6	17.6
<i>PapG III</i>	2	5.8
<i>sfa/focDE</i>	18	52.9
<i>traT</i>	2	5.8

In *E. faecalis*, the *efa* gene was detected in all isolates (100%), followed by *ebpA* (88.8%), *ebpB* (77.7%), *ebpC* (61.1%), *esp* (58.3%), *fsrA* (50.0%), *fsrB* (44.4%), *cylS* (38.8%), *cylL* (33.3%), and *fsrC* (27.7%), as shown in Table 10.

Table 10. Frequency of virulence genes detected in *E. faecalis*.

Gene	Positive isolates (n)	Percentage (%)
<i>efa</i>	36	100
<i>ebpA</i>	32	88.8
<i>ebpB</i>	28	77.7
<i>ebpC</i>	22	61.1
<i>esp</i>	21	58.3
<i>cylL</i>	12	33.3
<i>cylS</i>	14	38.8
<i>fsrA</i>	18	50.0

<i>fsrB</i>	16	44.4
<i>fsrC</i>	10	27.7

Among *S. aureus* isolates, *clfA* was the most frequently detected virulence gene (16.8%), followed by *agrIII* (8.9%), *etb* (6.7%), *agrII* (5.6%), *agrI* (4.4%), *eta* (3.3%), and *TSST-1* (2.4%). No *coa*, *X*-region or IgG-binding region genes were found. The corresponding frequencies for *S. aureus* and *P. aeruginosa* are presented in Table 11 and Table 12.

Among *P. aeruginosa* isolates, *exoS* was the predominant virulence gene (80.5%), followed by *plcH* (50.0%), *plcN* (47.2%), *exoY* (36.1%), *pilA* (30.5%), *pvdA* (27.7%), *phzII* (22.2%), *pilB* (22.2%), *toxA* (19.4%), *algU* (19.4%), *exoT* (19.4%), *phzM* (11.1%), *lasB* (11.1%), *algD* (8.3%), *apr* (8.3%), *nanI* (8.3%), *exoU* (5.5%), *phzI* (5.5%), *lasA* (5.5%), *phzH* (2.7%), and *phzS* (2.7%).

Table 11. Frequency of virulence genes detected in *S. aureus*.

Gene	Positive isolates (n)	Percentage (%)
<i>coa</i>	0	0
<i>clfA</i>	15	16.8
<i>X-region</i>	0	0
<i>IgG-binding region</i>	0	0
<i>TSST-1</i>	2	2.4
<i>eta</i>	3	3.3
<i>etb</i>	6	6.7
<i>agrI</i>	4	4.4
<i>agrII</i>	5	5.6

<i>agrIII</i>	8	8.9
---------------	---	-----

Table 12. Frequency of virulence genes detected in *P. aeruginosa*.

Gene	Positive isolates (n)	Percentage (%)
<i>exoT</i>	7	19.4
<i>exoY</i>	13	36.1
<i>exoS</i>	29	80.5
<i>exoU</i>	2	5.5
<i>phzH</i>	1	2.7
<i>apr</i>	3	8.3
<i>phzII</i>	8	22.2
<i>phzM</i>	4	11.1
<i>phzI</i>	2	5.5
<i>phzS</i>	1	2.7
<i>lasA</i>	2	5.5
<i>lasB</i>	4	11.1
<i>pvdA</i>	10	27.7
<i>pilA</i>	11	30.5
<i>pilB</i>	8	22.2
<i>toxA</i>	7	19.4
<i>algD</i>	3	8.3
<i>algU</i>	7	19.4
<i>plcH</i>	18	50.0

<i>plcN</i>	17	47.2
<i>nanI</i>	3	8.3

Representative agarose gel electrophoresis of PCR amplification products of selected *P. aeruginosa* virulence genes is shown in **Figure 1**.

Figure 1: Representative agarose gel electrophoresis of PCR amplification products of selected *P. aeruginosa* virulence genes.




Figure 1. Representative agarose gel electrophoresis of PCR amplification products of selected *P. aeruginosa* virulence genes. PCR products were separated by agarose gel electrophoresis and visualized under UV illumination after staining. M indicates the DNA molecular weight marker. The expected amplicon sizes of the amplified products (284, 408, 608, and 1159 bp) are indicated above the corresponding bands. This figure represents successful amplification of representative virulence genes analyzed in the present study.

Discussion

As part of the present study, the antibiotic-resistant bacteria and virulence determinants in municipal wastewater were monitored. *S. aureus* (45.6%), *E. faecalis* (18.5%), *P. aeruginosa* (18.5%), and *A. baumannii* (17.4%) were the most common of the 195 isolates. 45.6% (89) of the isolates were capable of producing ESBL, highlighting wastewater as an important reservoir and dissemination route for AMR. ESBL production was highest in *P. aeruginosa* (83.3%), followed by *S. aureus* (66.3%), *A. baumannii* (58.8%) and *E. faecalis* (25.0%). The high rate in *P. aeruginosa* may be due to the ability to acquire and horizontally transfer resistance determinants and intrinsic resistance (31), and due to the environmental adaptability of this species.

The most common ESBL gene detected was BLACTX-M, which was found in 36.1% of *P. aeruginosa* and 29.4% of *A. baumannii*, which is in concert with the global predominance of BLACTX-M as an environmental ESBL (32). BLASHV and BLATEM were detected in *A. baumannii* (23.5% and 5.8%, respectively) and BLACTX-M and BLATEM in *P. aeruginosa* (36.1% and 27.7%, respectively). Several ESBL genes can be detected, suggesting plasmid- or chromosome-mediated acquisition and horizontal transfer and heightened resistance (33, 34).

In the case of *A. baumannii*, *fimH* was the most common virulence gene (82.3%), which has been associated with adhesion and biofilm formation (35). Other adhesion and iron-acquisition mechanisms, *afa/draBC* and *fyuA*, were found in 55.8% and 52.9%, respectively, of the isolates (36). In contrast, *csgA*, *iutA*, PAI, and *ibeA* were not present, which may indicate different environmental genetic backgrounds.

All *E. faecalis* isolates (100%) were found to carry *efa*, and *ebpA*, *ebpB* and *ebpC* were detected in 88.8%, 77.7% and 61.1% of the isolates, respectively (37). The presence of the *esp* gene was found in 58.3%, the *cylS* gene in 38.8%, and the *cylL* gene in 33.3%. Of these isolates, 50.0% and

44.4% contained *fsr* quorum-sensing genes, indicating that there are multiple virulence mechanisms in place (38).

S. aureus had lower rates of virulence genes. *clfA* was the most common (16.8%), followed by *agrIII* at 2.4–8.9%, *etb*, *agrII*, *agrI*, *eta*, and TSST-1. The lack of *coa*, X-region, and IgG-binding region genes, and the rarity of *agr* locus occurrence, may suggest that the bacteria in the environment have different genetic traits from highly virulent clinical isolates.

P. aeruginosa was the most virulent with the highest number of virulence determinants. In 80.5% of the occurrences, *exoS* was detected, indicating the high pathogenicity potential of this bacterium (39). The *plcH* and *plcN* genes were found in 50.0% of samples and 47.2% of samples, respectively, and *exoY* was found in 36.1% and *exoU* in only 5.5% of samples. Phenazine-associated genes occurred at 22.2% (*phzII*), 11.1% (*phzM*), 5.5% (*phzI*), and 2.7% (*phzH*), while *pvdA* occurred in 27.7%. Pilus genes of type IV, *pilA* and *pilB*, were found in 30.5% and 22.2%, respectively. The *algU* and *algD* genes were present in 19.4% and 8.3%, respectively, suggesting alginate production ability and biofilm formation ability.

Overall, the results indicate the potential of WWTPs as reservoirs, amplification sites, and dissemination pathways for ARB and virulence determinants. The ESBL rate in influent (45.6%) is higher than those typically seen in community-acquired infections and is indicative of high selection pressure in wastewater (40). AMR genes may be present in combination with multiple virulence factors, which further enhances the potential public health threat via contaminated water, recreation and irrigation (41).

The high prevalence of BLACTX-M, BLASHV and BLATEM is consistent with other wastewater studies which found these three too often coexist on plasmids (42). Plasmid-mediated transfer can aid in transfer to other species of bacteria, and combinations of virulence determinants can enhance

pathogenicity, as well as increase the potential for multidrug resistance (43). The wastewater resistome and mobilome need to be characterized using metagenomic and functional metagenomic approaches, and whole-genome sequencing could help to clarify the resistance plasmids, insertion sequences, and genomic islands and horizontal gene transfer mechanisms (44, 45).

Conventional treatment may be ineffective at removing the resistant bacteria and resistance genes, as *P. aeruginosa* and *A. baumannii* are prevalent and ESBL-producing (ESBL-PA and ESBL-AB, respectively) (41). Therefore, advanced oxidation, membrane filtration, and improved disinfection may minimize the release of advanced oxidation products into receiving waters (46). The high prevalence of *efa* in *E. faecalis* and the presence of adhesion- and biofilm-formation-related genes are also alarming, as they enable biofilm development that further enhances AMR and antimicrobial persistence (47). The coexistence of resistance and virulence factors can raise the threat for vulnerable groups (48).

A One Health approach identifies wastewater as a nexus of human, animal, and environmental sources of AMR, and wastewater can be an early warning system for emerging resistance (42). Culture-dependent and culture-independent approaches to systematic surveillance may help to detect resistance mechanisms early and provide public health evidence-based interventions. This diversity of ESBL combinations and virulence profiles indicates that genetic exchange within wastewater is still occurring. The occurrence of BLACTX-M-associated resistance in both *P. aeruginosa* and *A. baumannii* may be due to plasmid-mediated acquisition and horizontal transfer from environmental bacterial flora (49). Also, ESBL-producing bacteria in sewage sludge are an indication of adaptation to sewage and possibly environmental dissemination (32).

Conclusion

The present study identified a high frequency of phenotypic ESBL-positive results among the recovered bacterial isolates from municipal wastewater and urban wastewater; a variety of virulence genes associated with ESBL-producing bacteria were recovered. Wastewater has emerged as a significant environmental source of AMR, especially for BLACTX-M, and also of bacterial pathogenicity, with multiple virulence determinants commonly found in the bacteria isolated from the wastewater samples of *P. aeruginosa*, *A. baumannii*, *E. faecalis* and *S. aureus*. The results highlight the critical need to improve wastewater treatment technologies, broaden routine environmental monitoring of AMRs, and develop comprehensive public health approaches to reduce the environmental spread of antimicrobial-resistant bacteria. Long-term longitudinal surveillance of AMR and functional studies of resistance plasmids and virulence determinants, combined with whole-genome sequencing, will generate more information on the ecology and evolution of AMR and help inform effective intervention strategies to reduce AMR dissemination.

Conflict of Interest Disclosure

The authors declare no conflict of interest.

Ethical Approval

Prior to conducting the study, all necessary approvals were obtained from the Ethics Committee of Islamic Azad University, Shahrekord Branch. The Ethics Committee of Islamic Azad University, Shahrekord Branch reviewed and approved the research protocol (IR.IAU.SHK.REC.1403.044).

Artificial Intelligence (AI) Disclosure Statement

The authors declare that no Gen AI was used in the creation of this manuscript.

Funding/Support

This study was funded by Islamic Azad University, Shahrekord Branch, Iran.

Acknowledgements

This article was derived from a research project approved by the Research and Technology Deputy of the Islamic Azad University, Shahrekord Branch. Hereby,

Authors Contribution

Author contributions should be clarified based on the following items:

Conceptualization: Hamid Reza Mardani and Hassan Momtaz

Data curation: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Formal Analysis: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Funding acquisition: Hassan Momtaz

Investigation: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Methodology: Hamid Reza Mardani and Rahman Abdizadeh

Project administration: Hassan Momtaz

Resources: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Software: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Supervision: Hassan Momtaz

Validation: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Visualization: Hamid Reza Mardani and Hassan Momtaz

Writing – original draft: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

Writing – review & editing: Hamid Reza Mardani, Rahman Abdizadeh and Hassan Momtaz

References

1. Mukherjee M, Laird E, Gentry TJ, Brooks JP, Karthikeyan R. Increased antimicrobial and multidrug resistance downstream of wastewater treatment plants in an urban watershed. *Front Microbiol.* 2021;12:657353. doi: [org/10.3389/fmicb.2021.657353](https://doi.org/10.3389/fmicb.2021.657353)
2. Hubeny J, Korzeniewska E, Buta-Hubeny M, Zieliński W, Rolbiecki D, Harnisz M. Characterization of carbapenem resistance in environmental samples and *Acinetobacter* spp. isolates from wastewater and river water in Poland. *Sci Total Environ.* 2022;822:153437. doi: [10.1016/j.scitotenv.2022.153437](https://doi.org/10.1016/j.scitotenv.2022.153437)
3. Chen L, Jiao J, Liu S, Liu L, Liu P. Mapping the global, regional, and national burden of diarrheal diseases attributable to unsafe water. *Front Public Health.* 2023;11:1302748. doi: [org/10.3389/fpubh.2023.1302748](https://doi.org/10.3389/fpubh.2023.1302748)
4. Kow CS, Ramachandram DS, Hasan SS. The effect of curcumin on the risk of mortality in patients with COVID-19: A systematic review and meta-analysis of randomized trials. *Phytother Res.* 2022;36(9):3365. doi: [10.1002/ptr.7468](https://doi.org/10.1002/ptr.7468)
5. Rodriguez-Mozaz S, Chamorro S, Marti E, Huerta B, Gros M, Sànchez-Melsió A, et al. Occurrence of antibiotics and antibiotic resistance genes in hospital and urban wastewaters and their impact on the receiving river. *Water Res.* 2015;69:234-42. doi: [10.1016/j.watres.2014.11.021](https://doi.org/10.1016/j.watres.2014.11.021)

6. Willett JL, Robertson EB, Dunny GM. The phosphatase Bph and peptidyl-prolyl isomerase PrsA are required for gelatinase expression and activity in *Enterococcus faecalis*. *J Bacteriol.* 2022;204(7):e00129-22. doi: [10.1128/jb.00129-22](https://doi.org/10.1128/jb.00129-22)
7. Hirvonen A, Comero S, Tavazzi S, Mariani G, Cacciatori C, Maffettone R, et al. “Encyclopaedia Cloacae”—Mapping Wastewaters from Pathogen A to Z. *Microorganisms.* 2025;13(8):1900. doi: [org/10.3390/microorganisms13081900](https://doi.org/10.3390/microorganisms13081900)
8. Mthembu TP, Zishiri OT, El Zowalaty ME. Detection and molecular identification of *Salmonella* virulence genes in livestock production systems in South Africa. *Pathogens.* 2019;8(3):124. doi: [10.3390/pathogens8030124](https://doi.org/10.3390/pathogens8030124)
9. Bozorgomid A, Chegine Lorestani R, Rostamian M, Nemati Zargarani F, Shahvaisy-Zadeh Z, Akya A. Antibiotic resistance, virulence factors, and phylogenetic groups of *Escherichia coli* isolated from hospital wastewater: A case study in the west of Iran. *Environ Health Eng Manag.* 2023;10(2):131-9. doi: [10.34172/EHEM.2023.15](https://doi.org/10.34172/EHEM.2023.15)
10. Drane K, Sheehan M, Whelan A, Ariel E, Kinobe R. The role of wastewater treatment plants in dissemination of antibiotic resistance: source, measurement, removal and risk assessment. *Antibiotics (Basel).* 2024;13(7):668. doi: [org/10.3390/antibiotics13070668](https://doi.org/10.3390/antibiotics13070668)
11. Fadaei A, Sadeghi M. Evaluation and assessment of drinking water quality in Shahrekord, Iran. *Resources and Environment.* 2014;4(3):168-72. doi: api.semanticscholar.org/CorpusID:92211970

12. Chagas T, Seki L, Cury J, Oliveira J, Dávila A, Silva D, et al. Multiresistance, beta-lactamase-encoding genes and bacterial diversity in hospital wastewater in Rio de Janeiro, Brazil. *J Appl Microbiol.* 2011;111(3):572-81. doi: [10.1111/j.1365-2672.2011.05072.x](https://doi.org/10.1111/j.1365-2672.2011.05072.x)
13. Silvester R, Woodhall N, Nurmi W, Muziasari W, Cross G, Malham SK, et al. High-throughput qPCR profiling of antimicrobial resistance genes and bacterial loads in wastewater and receiving environments. *Environ Pollut.* 2025;373:126096. doi: [10.1016/j.envpol.2025.126096](https://doi.org/10.1016/j.envpol.2025.126096)
14. Association APH, Association AWW, Federation WPC, Federation WE. Standard methods for the examination of water and wastewater: American Public Health Association.; 1917. ISBN: 978-0-87553-299-8
15. Guo Z, Hatakeyama T, Yoshimura C, Wang T, Hatano Y. Basic influent sewage quality reflects sewershed characteristics in Tokyo city. *J Water Health.* 2022;20(6):972-84. doi: [10.2166/wh.2022.022](https://doi.org/10.2166/wh.2022.022)
16. Schuetz AN, Ferrell A, Hindler JA, Humphries R, Bobenchik AM. Overview of changes in the clinical and laboratory standards institute performance standards for antimicrobial susceptibility testing: M100 32nd and 33rd editions. *J Clin Microbiol.* 2025;63(9):e01623-23. doi: [10.1128/jcm.01623-23](https://doi.org/10.1128/jcm.01623-23)
17. Diefenbach RJ, Lee JH, Kefford RF, Rizos H. Evaluation of commercial kits for purification of circulating free DNA. *Cancer Genet.* 2018;228:21-7. doi: [10.1016/j.cancergen.2018.08.005](https://doi.org/10.1016/j.cancergen.2018.08.005)

18. Neamah AA. Molecular Detection of virulence factor genes in *Pseudomonas aeruginosa* isolated from human and animals in Diwaniya province. *Kufa J Vet Med Sci.* 2017;8(1):218-30. doi: [org/10.36326/kjvs/2017/v8i14300](https://doi.org/10.36326/kjvs/2017/v8i14300)
19. Ranjbar R, Naghoni A, Afshar D, Nikkhahi F, Mohammadi M. Rapid molecular approach for simultaneous detection of *Salmonella* spp., *Shigella* spp., and *Vibrio cholera*. *Osong Public Health Res Perspect.* 2016;7(6):373-7. doi: 10.1016/j.phrp.2016.10.002
20. Melati R, Nurjanah S, Rahayu W. Desain Primer Gen Virulensi *invA* untuk Identifikasi dan Sekuensing *Salmonella* pada Sampel Karkas Ayam. *J Ilmu Produksi Teknol Hasil Peternakan.* 2022;10(2):91-7. doi: [org/10.29244/jipthp.10.2.91-97](https://doi.org/10.29244/jipthp.10.2.91-97)
21. Ngamwongsatit N, Chaturongakul S, Aunpad R. Development and validation of an efficient multiplex PCR assay for simultaneous detection of six common foodborne pathogens and hygiene indicators. *Foodborne Pathog Dis.* 2023;20(6):222-9. doi: 10.1089/fpd.2022.0062.
22. AL-Hassani MAN, Abdul-Lateef LA. Molecular detection of some virulence genes of *Enterococcus faecalis* isolated from different clinical samples in Babylon province/Iraq. *Int J Health Sci.* 2022;6(S4):2377-91. doi: [org/10.53730/ijhs.v6nS4.7371](https://doi.org/10.53730/ijhs.v6nS4.7371)
23. Liu Y, Cao Y, Wang T, Dong Q, Li J, Niu C. Detection of 12 common food-borne bacterial pathogens by TaqMan real-time PCR using a single set of reaction conditions. *Front Microbiol.* 2019;10:222. doi: 10.3389/fmicb.2019.00222.
24. Galeb S, Zaki ME, Shrief R, Hassan R, Rizk M. Diagnostic value of multiplex polymerase chain reaction in detection of *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Stenotrophomonas maltophilia* from Sepsis in Pediatrics. *Recent Pat Biotechnol.* 2021;15(3):195-203. doi: 10.2174/1872208315666210719104623

25. Wahdan A, Ezzat M, Emam A, Husseiny WA, Abou El-Atta M, Abd-Allah EM, Meligy AMA, Abdel-Raheem SM, Almathen F, Al-Shami S, et al. One health monitoring of resistant *Pseudomonas aeruginosa* in aquatic, poultry, and human sources: virulence traits and *bla_{shv}* gene tracking. *Pathogens*. 2025; 14(10):983. doi: [org/10.3390/pathogens14100983](https://doi.org/10.3390/pathogens14100983)
26. Ejaz H. Dissemination of SHV, TEM and CTX-M genotypes in pseudomonas aeruginosa: a pre-eminent reason for therapeutic failure in pediatrics. *Ann Clin Lab Sci*. 2020;50(6):797-805. PMID: 33334796
27. Hetta HF, Al-Kadmy IM, Khazaaal SS, Abbas S, Suhail A, El-Mokhtar MA, et al. Antibiofilm and antivirulence potential of silver nanoparticles against multidrug-resistant *Acinetobacter baumannii*. *Sci Rep*. 2021;11(1):10751. doi: 10.1038/s41598-021-90208-4
28. Elal Mus T, Cetinkaya F, Cibik R, Soyutemiz GE, Simsek H, Coplu N. Pathogenicity determinants and antibiotic resistance profiles of enterococci from foods of animal origin in Turkey. *Acta Vet Hung*. 2017;65(4):461-74. doi: 10.1556/004.2017.044
29. Maleki DT, Ghalavand Z, Laabei M, Nikmanesh B, Hourri H, Kodori M, et al. Molecular analysis of accessory gene regulator functionality and virulence genes in *Staphylococcus aureus* derived from pediatric wound infections. *Infection, Genetics and Evolution*. 2019;73:255-60. doi: 10.1016/j.meegid.2019.05.013
30. Ertugrul BM, Oryasin E, Lipsky BA, Willke A, Bozdogan B. Virulence genes *fliC*, *toxA* and *phzS* are common among *Pseudomonas aeruginosa* isolates from diabetic foot infections. *Infect Dis (Lond)*. 2018;50(4):273-9. doi: 10.1080/23744235.2017.1393839

31. Clarke LM, O'Brien JW, Murray AK, Gaze WH, Thomas KV. A review of wastewater-based epidemiology for antimicrobial resistance surveillance. *J Environ Expo Assess*. 2024;3(1). doi: [10.20517/jeea.2023.29](https://doi.org/10.20517/jeea.2023.29)
32. Lee C, Lee JH, Park M, Park KS, Bae IK, Kim YB, et al. Biology of *Acinetobacter baumannii*: Pathogenesis, Antibiotic Resistance Mechanisms, and Prospective Treatment Options. *Front Cell Infect Microbiol*. 2017;7. doi: [10.3389/fcimb.2017.00055](https://doi.org/10.3389/fcimb.2017.00055)
33. Yehya A, Ezzeddine Z, Chakkour M, Dhaini Z, Bou Saba MS, Bou Saba AS, et al. The intricacies of *Acinetobacter baumannii*: a multifaceted comprehensive review of a multidrug-resistant pathogen and its clinical significance and implications. *Front Microbiol*. 2025;16. doi: [10.3389/fmicb.2025.1565965](https://doi.org/10.3389/fmicb.2025.1565965)
34. Sakalauskienė GV, Malcienė L, Stankevičius E, Radzevičienė A. Unseen Enemy: Mechanisms of Multidrug Antimicrobial Resistance in Gram-Negative ESKAPE Pathogens. *Antibiotics (Basel)*. 2025;14(1). doi: [10.3390/antibiotics14010063](https://doi.org/10.3390/antibiotics14010063)
35. Yang S, Meng X, Zhen Y, Baima Q, Wang Y, Jiang X, et al. Strategies and mechanisms targeting *Enterococcus faecalis* biofilms associated with endodontic infections: a comprehensive review. *Front Cell Infect Microbiol*. 2024;14. doi: [10.3389/fcimb.2024.1433313](https://doi.org/10.3389/fcimb.2024.1433313)
36. Li P, Zhang S, Wang J, Al-Shamiri MM, Han B, Chen Y, et al. Uncovering the Secretion Systems of *Acinetobacter baumannii*: Structures and Functions in Pathogenicity and Antibiotic Resistance. *Antibiotics (Basel)*. 2023;12(2). doi: [10.3390/antibiotics12020195](https://doi.org/10.3390/antibiotics12020195)
37. Barros V, Rocha BS, Rhoden J, Stein JF, Picoli SU, Rigotto C. Detection of metallo-beta-lactamase-producing genes *bla*SPM and *bla*NDM in *Pseudomonas aeruginosa* isolated from wastewater in Southern Brazil. *J Water Health*. 2024;22(4):689-700. doi: [10.2166/wh.2024.305](https://doi.org/10.2166/wh.2024.305)

38. Pandey RP, Yousef AF, Alsafar H, Hasan SW. Surveillance, distribution, and treatment methods of antimicrobial resistance in water: A review. *Sci Total Environ.* 2023;890:164360. doi: 10.1016/j.scitotenv.2023.164360
39. Kumar G, Balakrishna K, Mukhopadhyay C, Kalwaje Eshwara V. Characterization and comparative analysis of antimicrobial resistance in *Escherichia coli* from hospital and municipal wastewater treatment plants. *J Water Health.* 2024;22(12):2276-88. doi: 10.2166/wh.2024.126
40. Waseem H, Poorasgari E, Örmeci B. Tracking antibiotic resistance and respiratory viruses in municipal wastewater: Site-specific dynamics from two wastewater treatment plants and an upstream Sewershed community. *Int J Hyg Environ Health.* 2026;274:114767. doi: 10.1016/j.ijheh.2026.114767
41. Knight ME, Webster G, Perry WB, Baldwin A, Rushton L, Pass DA, et al. National-scale antimicrobial resistance surveillance in wastewater: A comparative analysis of HT qPCR and metagenomic approaches. *Water Res.* 2024;262:121989. doi: 10.1016/j.watres.2024.121989
42. Silvester R, Webster G, Perry WB, Farkas K, Rushton L, Craine N, et al. Genome-resolved metagenomics uncovers antimicrobial resistance gene carriers in hospital and municipal wastewater environments. *Sci Total Environ.* 2025;180607. doi: 10.1016/j.scitotenv.2025.180607
43. Taylor W, Bohm K, Dyet K, Weaver L, Pattis I. Comparative analysis of qPCR and metagenomics for detecting antimicrobial resistance in wastewater: a case study. *BMC Res Notes.* 2025;18(1). doi: 10.1186/s13104-024-07027-9
44. Roy S, Chowdhury G, Mukhopadhyay AK, Dutta S, Basu S. Convergence of Biofilm Formation and Antibiotic Resistance in *Acinetobacter baumannii* Infection. *Front Med (Lausanne).* 2022;9. doi: 10.3389/fmed.2022.793615

45. Islam MM, Jung DE, Shin WS, Oh MH. Colistin Resistance Mechanism and Management Strategies of Colistin-Resistant *Acinetobacter baumannii* Infections. *Pathogens*. 2024;13(12). doi: 10.3390/pathogens13121049
46. Punch R, Azani R, Ellison C, Majury A, Hynds PD, Payne SJ, et al. The surveillance of antimicrobial resistance in wastewater from a one health perspective: A global scoping and temporal review (2014–2024). *One Health*. 2025;21:101139. doi: 10.1016/j.onehlt.2025.101139
47. Bydalek F, Webster G, Barden R, Weightman AJ, Kasprzyk-Hordern B, Wenk J. Microplastic biofilm, associated pathogen and antimicrobial resistance dynamics through a wastewater treatment process incorporating a constructed wetland. *Water Res*. 2023;235, 119936. doi: org/10.1016/j.watres.2023.119936
48. Raven KE, Ludden C, Gouliouris T, Blane B, Naydenova P, Brown NM, et al. Genomic surveillance of *Escherichia coli* in municipal wastewater treatment plants as an indicator of clinically relevant pathogens and their resistance genes. *Microb Genom*. 2019;5(5): e000267. doi: 10.1099/mgen.0.000267
49. Novović K, Jovčić B. Colistin Resistance in *Acinetobacter baumannii*: Molecular Mechanisms and Epidemiology. *Antibiotics (Basel)*. 2023;12(3). doi: 10.3390/antibiotics12030516